


# LONG-TAILED JAEGER

*Stercorarius longicaudus*



## Regional Occurrence (BCRs 14, 30)

- Distribution: pelagic in region
- Habitats used: open or alpine tundra, flats with sparse vegetation (breeding); pelagic, along seacoasts and on inland waters (non-breeding)

NAWCP BD=1; NAWCP ND=1

## Regional Abundance

- 1980s population size: 100-1,000nb
- Historical population size: unknown
- % change in regional population: unknown

NAWCP PS=2; NAWCP PT=3

## Regional Threats

- Likely subject to environmental pollutants in oceans
- Susceptible to predation, especially by arctic foxes
- Success of fledglings is dependent on conditions at sea

NAWCP TB=3; NAWCP TN=2

**Conservation Status:** Abundance (b=breeding, nb=nonbreeding, t=total individuals) and Conservation Rank

Global	North America	BCR 14	BCR 30	Mid-Atlantic	S. New England	Gulf of Maine	Maritimes
?	>150,000t	?nb	?nb	?nb	?nb	?nb	?nb
IUCN- Not Evaluated	NAWCP- Low Concern	Low (PI=?)	Low (PI=?)	NJ- SZN; DE-SZN MD-SZN; VA-SN	NY-S?N; CT-no info RI- no info; MA, VT-no info	MA-no info; NH- no info ME-SZN; NB- no info, NS-SZN	QC-S3; NB- no info PE-no info; NS-SZN

**Conservation Needs:** *Sustainability:* 1.4 young fledged / pair; *Foraging Habitat:* males on outer part of territory or sometimes at sea; females within few hundred meters; non-breeding not usually less than 35 km from coast; *Issues:* Management has not occurred in natural populations, and the species generally does not fair well in captivity.; *Population Goals:* ; *Habitat Goals:*